



Australian Government
Department of Industry,
Innovation and Science

National Measurement Institute



REFERENCE MATERIAL CATALOGUE

WADA & PCC funded Reference Materials Catalogue

NMI reference materials

available globally through our international partners:



Cerilliant[®]
Analytical Reference Standards

Cerilliant

www.cerilliant.com



LGC Standards

www.lgcstandards.com

MERCK

Merck

www.sigmaaldrich.com

Add more body copy here if you want or delete.

	NMI reference	Unit of issue (mg)	Cost (A\$)	
			Domestic (excl GST)	International
STEROIDS AND STEROID METABOLITE				
1-Androstendione				
1-Androstendione (5 α -Androst-1-en-3,17-dione) *	D845	1	207	290
Danazol metabolites				
2 α -Hydroxymethylethisterone †	D920b	1	110	154
Finasteride metabolites				
Carboxy finasteride †	S045	1	110	154
4-Hydroxy steroids				
4-Hydroxy-estr-4-en-3,17-dione †	S043	1 (ampoule)	110	154
17α-Methylclostebol metabolite				
4-Chloro-17 α -methyl-androst-4-ene-3 α , 17 β -diol †	S044	1 (ampoule)	110	154
Methyldienolone				
Methyldienolone †	D916b	1 (ampoule)	110	154
7α-Methylnandrolone and metabolites				
7 α -Methylnandrolone *	S048	1 (ampoule)	110	154
7 α -Methyl-5 β -estran-3 α -ol-17-one (major metabolite) *	S047	1 (ampoule)	110	154
7 α -Methyl-estr-4-ene-3 α -ol-17-one (minor metabolite) *	S050	1 (ampoule)	110	154
Nandrolone metabolites				
19-Noretiocholanolone sulfate (Na salt)*	D849	1	110	154
Epinandrolone sulfate† TEA salt	D783b	5	110	154
Norbolethone and metabolites				
Norbolethone† (13 β ,17 α -Diethyl-gonan-4-ene-17 β -ol-3-one)	D825c	1 (ampoule)	110	110
13 β ,17 α -Diethyl-5 β -gonane-3 α , 17 β -diol (major metabolite) †	D818b	1 (ampoule)	110	154
13 β ,17 α -Diethyl-5 α -gonane-3 α , 17 β -diol (minor metabolite) †	D820b	1 (ampoule)	110	154
Testosterone metabolites (including internal standards)				
5 β -Androstane-3 α ,17 β -diol-3- β -glucuronic acid †	S003b	1	110	154
d4-5 α -Androstan-3 α ,17 β -diol-17-O- β -glucuronic acid *	S009	1	110	154
d4-5 α -Androstan-3 α ,17 β -diol-3-O- β -glucuronic acid *	S010	1	110	154
d3-5 β -Androstan-3 α ,17 β -diol-17-O- β -glucuronic acid *	S011	1	110	154
d5-5 β -Androstan-3 α ,17 β -diol-3-O- β -glucuronic acid *	S012	1	110	154
d4-Epitestosterone-17-O- β -glucuronic acid *	S023	1	110	154
d5-Etiocholanolone-3-O- β -glucuronide sodium salt *	S020	1	110	154
1-Testosterone*				
5 α -Androst-1-ene-3 α -ol-17-one* (Ampouled)	D832	0.2	110	154
Oral Turinabol metabolite				
6 β -Hydroxy-oral turinabol †	D615b	1 (ampoule)	110	154

	NMI reference	Unit of issue (mg)	Cost (A\$)	
			Domestic (excl GST)	International
Prohormones and metabolites				
3 α ,5-Cyclo-5 α -androstan-6 β -ol-17-one †	S039	1 (ampoule)	110	154
16 α -Hydroxyandrosterone *	D843	1 (ampoule)	110	154
6 β -Hydroxyetiocholanolone *	D867	1	110	154
4 β -Hydroxy DHEA *	D834	1 (ampoule)	110	154
7-Keto DHEA *	D833	10	110	154
7 α -Hydroxy DHEA *	D875	1	110	154
7 β -Hydroxy DHEA *	D865b	1	110	154
16 β -Hydroxy DHEA *	D844	1	110	154
3 α -Hydroxy-4-estren-17-one *	D873	1	110	154
3 β -Hydroxy-4-estren-17-one *	D866	1	110	154
Anti-inflammatory metabolites				
α -Hydroxycarprofen†	D1072	5	110	154
REV-ERB agonist SR9009 metabolites				
<i>N</i> -[(4-Chlorophenyl)methyl]-5-nitro-2-thiophenemethanamine hydrochloride†	D1066	1 (ampoule)	110	154
Ethyl <i>N</i> -(5-nitro-2-methylthiophene)-3-aminomethylpyrrolidine-1-carboxylate†	D1067	1 (ampoule)	110	154
Stimulants				
Formoterol fumarate†	D1065	25	110	154
Higenamine hydrochloride†	D1070	25	110	154
STERIODS AND STEROID METABOLITES				
Steroid carbon isotope ratio standard mixture reference material for GCC-IRMS				
Carbon isotope delta value reported for 5 α -androstan-3 β -ol acetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -32.00 \pm 0.09), 5 α -androstan-3 α -ol-17-one acetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -32.58 \pm 0.03), 5 β -androstan-3 α -ol-11,17-dione acetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -16.50 \pm 0.03), and 5 α -cholestane ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -24.90 \pm 0.05) †	CU-PCC 33-2	Ampoule	110	154
Carbon isotope delta value reported for 5 β -androstan-3 α -ol-17-one ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -28.75 \pm 0.01), 5 α -androstan-3 α -ol-17-one ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -26.74 \pm 0.03), 5 β -pregnan-3 α , 20 α -diol ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -18.65 \pm 0.03) †	CU-PCC 34-3	Ampoule	110	154
Carbon isotope delta value reported for 5 α -androstan-3 α -ol-17-one ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -27.09 \pm 0.07) †	CU-PCC 40-1	Ampoule	110	154
Carbon isotope delta value reported for 5 α -androstan-3 α -ol-17-one acetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -32.82 \pm 0.02) †	CU-PCC 41-1	Ampoule	110	154
Carbon isotope delta value reported for 5 α -cholestane ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -25.03 \pm 0.01) †	CU-PCC 42-1	Ampoule	110	154
Carbon isotope delta value reported for 5 α -androstan-3 α -ol-17-one acetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -32.73 \pm 0.06), 5 β -androstan-3 α , 17 β -diacetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -30.19 \pm 0.07), 5 α -cholestane ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -24.83 \pm 0.13), and 5 β -pregnan-3 α , 20 α -diacetate ($\delta^{13}\text{C}_{\text{VPDB}}$ / ‰ -21.16 \pm 0.08) †	CU-PCC 44-1	Ampoule	110	154
Steroid Matrix Reference Material				

	NMI reference	Unit of issue (mg)	Cost (A\$)	
			Domestic (excl GST)	International
19-Norandrosterone (221.4 ng/g) in 1,2-dimethoxyethane (1 mL) *	MX003	Ampoule	165	242
Carbon Isotope Delta Value of 19-norandrosterone in Water containing 20% Methanol 1 (mL) * $\delta^{13}\text{C}_{\text{VPDB}} / \text{‰} -29.7 \pm 0.8$	MX016	Ampoule	240	336
Mass fraction of testosterone metabolites in freeze dried human urine: 5 α -androstane-3 α -17 β -diol (41.2 \pm 1.8 ng/g), 5 β -androstane-3 α -17 β -diol (66.0 \pm 2.9 ng/g), androsterone 1652 \pm 29 ng/g), etiocholanolone (1359 \pm 36 ng/g), testosterone (88.1 \pm 4.2 ng/g), epitestosterone (21.9 \pm 1.0 ng/g), T/E mass ratio (4.03 \pm 0.26) *	MX017	Bottle	240	336
Carbon Isotope Delta Value ($\delta^{13}\text{C}_{\text{VPDB}} / \text{‰}$). Three ampoules containing dry steroid mixtures. The ampoules contain approximately 400 μg of each steroid with the exception of 16-androstenol supplied close to 280 μg . Vial 1: etiocholanolone (-27.94 \pm 0.24), androsterone (-27.79 \pm 0.21), 11-oxoetiocholanolone (-13.58 \pm 0.23), testosterone (-27.87 \pm 0.24), 11 β -hydroxyetiocholanolone -29.51 \pm 0.36) Vial 2: 5 β -androstane-3 α -17 β -diol (-29.86 \pm 0.16), 5 α androstane-3 α -17 β -diol (-31.14 \pm 0.24), pregnanediol (-16.79 \pm 0.42), epitestosterone (-30.17 \pm 0.36), 11 β -hydroxyandrosterone (-28.59 \pm 0.22) Vial 3: 16-androstenol (-30.96 \pm 0.37), dehydroepiandrosterone (-31.63 \pm 0.54), testosterone (-22.52 \pm 0.33) †	MX018	3 Ampoules	240	336

*Production funded by the World Anti-Doping Agency.

†Production funded by the Partnership for Clean Competition.

NOTES

1. Restrictions on Issue

Production of these materials is funded by the World Anti-Doping Agency. These materials are only available to WADA accredited laboratories, or to laboratories which have entered the WADA accreditation process. Non-WADA-accredited laboratories who wish to purchase these materials must seek approval from WADA and provide evidence of the approval to NMI.

2. Classes of Reference Materials available from NMI

2.1 NMI Reference Materials

Reference Materials (RMs) produced by NMI are prepared in accordance with our accreditation from NATA to ISO 17034:2016(E) *General requirements for the competence of reference materials producers*. RMs are supplied with product information sheets and SDS, and have demonstrated homogeneity and stability.

2.2 NMI Certified Reference Materials

Certified Reference Materials (CRMs) produced by NMI have established metrological traceability to the SI unit for mass (kg). All purity values, stated as a mass fraction, are reported with the associated uncertainty in the certificate of analysis.

3. Ampouled Materials

For each ampouled reference material the product information sheet / certificate of analysis will state the mass of analyte dispensed into each ampoule. For reference materials with certified reference material status the mass of analyte will be reported with an associated uncertainty.

4. GST charges (Australian customers only)

The prices quoted are EXCLUSIVE of Goods and Services Tax (GST). The current GST rate is 10%. Upon delivery of the requested reference materials NMI will issue valid tax invoices and, if required, adjustment notes as per the requirements of the GST legislation.

5. Delivery fee

Delivery and handling fees apply for all orders and vary with destination. Please check the applicable rates when ordering.

6. For further information

Customer Service Officer,
Chemical Reference Materials Team,
National Measurement Institute,
GPO Box 2013, Canberra, ACT 2601, Australia.

Phone: +61 2 9449 0191

Fax: +61 2 9449 0292

Email: chemref@measurement.gov.au

Web: www.measurement.gov.au/chemref

NMI reference materials available globally through our international partners:



Cerilliant

www.cerilliant.com



LGC Standards

www.lgcstandards.com



Merck

www.sigmaaldrich.com